
CYTOPLASM STIFFNESS CAN REVERSE NUCLEAR PLASTICITY IN LAMIN-DEFICIENT CELLS

Solenne Deveraux¹, Rachele Allena², Denis Aubry¹

¹ MSSMat, CentraleSupelec, Université Paris-Saclay, 92290 Châtenay-Malabry, France ² Arts et Metiers ParisTech, LBM/Institut de Biomecanique Humaine Georges Charpak, 75013 Paris, France



Comparison of the perfusion test results in the 1 µm-wide micro-channel for wild type and lamin deficient cells. Left: Resulting fluid force (vertical and horizontal components), Right: Nucleus deviatoric strain

- [1] Teresa Audesirk & Audesirk, Gerald. Biology life on earth (5th ed) (1999). Upper Saddle River, N.J Prentice Hall
- [2] 1st Place in 2003 Nikon Photomicrography Competition, Dr. Torsten Wittmann
- [3] Lämmermann, Tim, and Michael Sixt. "Mechanical Modes of 'Amoeboid' Cell Migration." Current Opinion in Cell Biology 21.5 (2009): 636–644. NCBI PubMed. Web.
- [4] Columbia University Medical Center (2009)
- [5] Pajerowski, J. David et al. "Physical Plasticity of the Nucleus in Stem Cell Differentiation." Proceedings of the National Academy of Sciences of the United States of America 104.40 (2007): 15619–15624. PubMed Central. Web.

